

# CHITINOZOA FROM THE ORDOVICIAN SYLVAN SHALE OF THE ARBUCKLE MOUNTAINS, OKLAHOMA

by W. A. M. JENKINS

**ABSTRACT.** Chitinozoa referable to nine genera and twelve species (five new) are recorded from the Sylvan Shale of the Arbuckle Mountains in southern Oklahoma. They provide for the first time a means of dating, biostratigraphically subdividing, and correlating the entire formation. Its chitinozoan fauna indicates that the Sylvan Shale is of Upper Ordovician age throughout and confirms with additional fossil evidence the age of the lower beds. No abrupt changes break the general continuity of the chitinozoan succession but on the basis of gradual changes in the composition of its chitinozoan fauna the Sylvan Shale may be divided into three biostratigraphical units. The fauna occurs throughout the Arbuckle Mountains and is recognizable in the subsurface of western Texas and in the Maquoketa Formation of eastern Iowa. It differs strikingly from the fauna in the underlying Viola and Fernvale Limestones, and from the faunas in the overlying formations of the Silurian-Devonian Hunton Group. The occurrence of reworked Sylvan chitinozoans in upper Sylvan strata indicates that during late Sylvan time uplift (hitherto unsuspected) took place, at least locally, in the southern mid-continent.


The fauna is described systematically. *Clathrochitina* Eisenack 1959 is used in its original sense; *Plectochitina* Cramer 1964 is a junior synonym. *Sagenachitina* gen. nov. is proposed for species, hitherto referred to *Clathrochitina*, whose basal margins support finely divided networks. *Acanthochitina rashidi*, *Ancyrochitina merga*, *Clathrochitina sylvanica*, *Cyathochitina agrestis*, and *Sphaerochitina lepta* are new species.

THIS account of chitinozoans from the Sylvan Shale is the second of two investigations directed primarily at establishing an Upper Ordovician chitinozoan reference section in the Arbuckle Mountains of southern Oklahoma. In a stratigraphical sense it is the continuation of an earlier study (Jenkins 1969) devoted to chitinozoans from the underlying Viola and Fernvale Limestones. An incidental objective was to provide, for the first time, a means of dating and correlating the entire formation.

The Sylvan Shale is the youngest Ordovician formation in the Arbuckle Mountains, and rests unconformably upon the slightly older Viola-Fernvale limestone succession. It was chosen for this investigation primarily because it contains large numbers of well-preserved chitinozoans at practically all stratigraphical levels, but several additional factors also favoured its selection as a standard for stratigraphical reference. In particular, it is a distinctive lithostratigraphical unit separated by regional unconformities and by its sharply contrasting lithology from thick carbonate sequences above and below. In the subsurface the Sylvan Shale extends across most of Oklahoma into north-western Texas and southern Kansas and has been widely used as a marker horizon in the subsurface mapping of an otherwise unbroken sequence of Ordovician to Devonian limestones.

The formation crops out widely within the Arbuckle Mountains, for which reason the samples were collected there. It is particularly well exposed and free from faulting in a narrow outcrop 14 miles (22 km.) long, running west-north-west from Ravia to the Washita River along the steeply dipping southern flank of the Sycamore Creek Anticline (text-fig. 1). Chitinozoans from this outcrop form the basis of the present study.

In this paper, for reasons presented elsewhere (Jenkins 1969, pp. 5, 6), the base of the *Nemagraptus gracilis* Zone (or its correlatives) is taken as the lower limit of the Upper Ordovician. Thus defined, the Upper Ordovician includes the basal beds of the North



TEXT-FIG. 1. Showing the outcrop (outlined) and structural features of the pre-Pennsylvanian rocks in the Arbuckle Mountains. The shaded portion of the index map of Oklahoma delineates the area shown. The upper left inset map (represented by the shaded rectangle in the chart) shows the outcrop (stippled) of the Sylvan Shale exposed in the Sycamore Creek Anticline, and the road from Baum (on U.S. Highway 177) to the section (S) which provided the rock samples for this study. Compiled from Ham, W. E. 1955, *Geology of the Arbuckle Mountain Region*, Guide Book III; Ham, W. E. *et al.*, 1954, *Geologic Map and Sections of the Arbuckle Mountains, Oklahoma*, Guide Book III. Both published by the Oklahoma Geological Survey.

American Black River Stage and their correlatives (Twenhofel *et al.* 1954; Berry 1960a, b; Kay 1960, p. 30) at the base of the Caradoc Series in Britain.

#### PREVIOUS RESEARCH

Previous publications concerned with chitinozoans from the Sylvan Shale are limited to two notes (Wilson 1958, Wilson and Hedlund 1964) and an abstract (Wilson and Hedlund 1962). The latter is from Hedlund's (1960) substantial account of chitinozoans, acritarchs, and scolecodonts, which was submitted as a thesis for the Master of Science degree to the University of Oklahoma. Although unpublished, Hedlund's work has remained for a decade the chief source of information about

organic-walled microfossils in the Sylvan Shale, and the only attempt to describe them systematically.

The possibility exists, however, that chitinozoans were encountered in fossil residues of the Sylvan Shale as early as 1930. A short publication by Thomas (1930) contains (p. 87) not only a very early (perhaps the earliest) reference to organic-walled microfossils in the Lower Palaeozoic of Oklahoma, but almost certainly a passing reference to chitinozoans. I have been unable to ascertain whether or not this paper preceded Eisenack's of the same year, in which chitinozoans (though not named as such) were introduced to the scientific literature. On account of this paper's apparent obscurity outside North America the following relevant passage is reproduced.

... I also have permission from Mr. S. W. Lowman, micropaleontologist for the Midcontinent Petroleum Corporation, to announce a discovery he has recently made concerning the Sylvan in Oklahoma and Kansas. He has found that after completely dissolving the shale in concentrated hydrochloric acid the residue will yield microfossils which are invisible upon ordinary microscopic examination. He discovered organisms similar to *Sporangites* found in the Woodford [Formation, which is Devonian-Mississippian]. One species is about as large as the disc-like *Sporangites huonense* and is found in the upper Sylvan from central Kansas to the Arbuckles. Another species about twice as large as the above has been found in the upper Sylvan on the outcrop and in wells near the Arbuckles. A few spindle-shaped forms have been found. There are reasons to believe that these are plant spores and there are also arguments in favor of the theory that they are graptolitic in nature (Thomas 1930, p. 87).

In all probability the 'spindle-shaped forms' are chitinozoans for, to my knowledge, no organic-walled microfossils in the Sylvan Shale other than certain chitinozoans (and possibly a few netromorph acritarchs) could be interpreted, however broadly, as spindle-shaped. It is interesting to learn from Thomas's paper that the speculation about the natural affinities of the Chitinozoa, which has continued to the present day, had already started before the group was formally established and named by Eisenack in 1931.

#### GENERAL STRATIGRAPHY

*Lithology and fauna.* Throughout the Arbuckle Mountains the lower part of the Sylvan Shale, up to c. 130 ft. (40 m.) thick, is a hard, splintery, very fissile, brown to dark grey shale containing graptolites which are particularly abundant near the base. By contrast, the upper part of the formation, up to 200 ft. (61 m.) thick, is mainly a soft, weakly fissile to concretionary, greenish-grey shale, and is apparently devoid of macrofossils. The topmost 30 ft. (9 m.) is a soft, massive, highly pyritic, light green claystone, which disaggregates completely in water and is, for this reason, only ever temporarily exposed. The Sylvan Shale is generally slightly calcareous, and may be highly calcareous locally, particularly in its upper 100 ft. (30 m). Dolomite beds a few inches thick occur irregularly throughout the formation, and in the lower part two beds of dense brown dolomite, each 1-3 ft. (0.30-0.91 m.) thick, extend laterally over a wide area. Outcrops of the Sylvan Shale are strikingly delineated by sharp changes in the topography, soil, and vegetation of the limestone country within which they are exposed.

The formation contains, in addition to one species of brachiopod (Cooper 1956, p. 244) and seven species of graptolites (Decker 1935, 1945; Ruedemann 1947, pp. 90-2, 100) in its lower part, a rich fauna of chitinozoans and acritarchs throughout. Generally,

however, other fossils are lacking, and the upper Sylvan in particular is seemingly devoid of macrofossils.

The lithology and fauna of the Sylvan Shale indicate that it was laid down in tranquil waters, and suggested to Ham (1969, p. 11) that it was deposited in deeper waters than all other pre-Mississippian formations in the Arbuckle Mountains. It is the only pre-Upper Devonian shale unit, moreover, to persist throughout the Arbuckle Geosyncline and to extend beyond it over much of the southern mid-continent.

*Pattern of deposition.* Within the Arbuckle Mountains the Sylvan Shale's broad pattern of deposition is similar to that of the Viola Limestone (Jenkins 1969, p. 3). The Sylvan sediments in the south-western Arbuckle Mountains were laid down in the rapidly subsiding basin of the Arbuckle Geosyncline, whereas those to the north and east of a line now approximately followed by the Reagan Fault (text-fig. 1) were deposited on a slowly subsiding geosynclinal shelf. This led to the accumulation of less sediment in the north-east than in the south-west and explains the general thinning of the formation north-eastward across the Arbuckle Mountains from a maximum of 325 ft. (99 m.) in the basin to 150–175 ft. (45–53 m.) on the shelf (Ham 1969, p. 10).

*Stratigraphical relations.* Over a large part of Oklahoma, including the Arbuckle Mountains, regional unconformities separate the Sylvan Shale from a thick sequence of older Ordovician limestones below, and a succession of Silurian and Devonian carbonate formations above. The Sylvan lies everywhere upon the Fernvale Limestone, and the sharp contact is well exposed in Sycamore Creek (text-fig. 1), map reference SW $\frac{1}{4}$  SE $\frac{1}{4}$  sec. 27, T. 3 S., R. 4 E., Johnston County, Oklahoma. The dips of the two formations appear to be the same, but close examination reveals that the top of the Fernvale Limestone is an erosion surface that had been extensively weathered before the shale was laid down. A 2–3 in. (5–8 cm.) thick band of oxidizing pyritic 'rubble', containing phosphatized pebbles, locally marks the base of the shale, as at Sycamore Creek, and (Ham, personal communication, May 1969) is particularly well developed as a pyritic phosphate conglomerate, containing dolomite pebbles, in the road-cut entrance to Rayford Quarry (text-fig. 1), map reference NW $\frac{1}{4}$  NE $\frac{1}{4}$  NE $\frac{1}{4}$  sec. 28, T. 1 S., R. 2 E., Murray County, Oklahoma. This and related late Ordovician unconformities are given some prominence, and discussed in a regional context, by Ham and Wilson (1967, pp. 350–2) in a comprehensive account of Palaeozoic tectonic disturbances in the central United States.

Almost everywhere within the Arbuckle Mountains the Sylvan Shale is unconformably overlain by Silurian or Lower Devonian limestones (Amsden 1960, panel III). These are extremely thin within the Sycamore Creek Anticline, however, and locally, as at Sycamore Creek, they have been entirely removed by mid-Devonian erosion; the black shales and bedded cherts of the Upper Devonian–Mississippian Woodford Formation lie directly upon the Sylvan Shale. The Woodford Formation, which is 350–560 ft. (107–171 m.) thick in the Arbuckle Mountains, has been dated by conodonts as ranging in age from Frasnian (Upper Devonian) to Kinderhookian (Lower Mississippian); the Mississippian part of the formation generally is less than 1 ft. (0.30 m.) thick, and exceptionally at least 10 ft. (3 m.) thick (Hass and Huddle 1965).

*Age and correlation.* The Sylvan Shale was dated Upper Ordovician by Decker (1935), on the basis of graptolites confined to the lower part of the formation. The greater part



of the Sylvan Shale has yielded no macrofossils, however, but was placed in the Ordovician on the basis of its association with the lower Sylvan. Although Decker's age determination for the lower graptolitic Sylvan has generally been accepted, the age of the overlying 'unfossiliferous' beds, at least 200 ft. (61 m.) thick in the south-western Arbuckle Mountains, has not hitherto been demonstrated.

*Collection and preparation of material.* Samples were collected at stratigraphical intervals of 20 ft. (6.10 m.) throughout the Sylvan Shale succession exposed in the bed and banks of Sycamore Creek (text-fig. 1), map reference SW $\frac{1}{4}$  SE $\frac{1}{4}$  sec. 27, T. 3 S., R. 4 E., Johnston County, Oklahoma, commencing 1 ft. above the Fernvale Limestone-Sylvan Shale contact. The latter serves as an easily recognizable reference datum, and samples are numbered according to their original stratigraphical positions (measured in feet) above it. For example, sample Sy280 is from a stratigraphical level 280 ft. above the contact, the prefix 'Sy' indicating that it is from Sycamore Creek. These samples are briefly described in the appendix.

The collection of regularly spaced samples was made easy at Sycamore Creek by the steep inclination of the beds and the absence of faulting. The relatively resistant shales in the lower part of the formation are well exposed in the bed and steep banks of the creek, and samples of these were collected without difficulty. To obtain samples of the softer shales and claystones near the top of the formation, however, it was necessary to dig through as much as 2 ft. (0.6 m.) of Recent alluvium. The total thickness of the Sylvan Shale at Sycamore Creek, from its sharp contact with the Fernvale limestone below, to its equally sharp contact with the Woodford Formation above, was carefully measured by Mr. J. A. Turnbull and myself, and found to be 305 ft. (93 m.), slightly less than Decker's (1935, p. 698) figure of 320 ft. (97.5 m.).

In order to determine how much the character of the Sylvan chitinozoan fauna varies regionally several samples were collected at widely scattered localities (q.v., p. 283) throughout the Arbuckle Mountains. A few figured specimens are from outcrops along U.S. Highway 77 about 4 miles (6.4 km.) south of Davis (text-fig. 1), map reference sec. 30, T. 1 S., R. 2 E., Murray County, Oklahoma. Their reference numbers are prefixed 'Da'.

The chitinozoans were prepared for microscopical examination according to the method outlined by Jenkins (1967, pp. 439-41), and furnished with reference numbers in the manner described for the Viola and Fernvale specimens of an earlier study (Jenkins 1969, p. 7). Most of the preparations, including all the type material, and portions of each rock sample, are housed in the Micropalaeontology Laboratory, Department of Geology, The University, Sheffield, England.

## SYSTEMATIC PALAEOLOGY

### Order CHITINOZOA Eisenack 1931

#### Genus ACANTHOCHITINA Eisenack 1931 emend. Jenkins 1967

*Type species.* *Acanthochitina barbata* Eisenack 1931 (by original designation), Ordovician, Baltic.

#### *Acanthochitina rashidi* sp. nov.

Plate 47, fig. 20; Plate 48, figs. 1, 2; text-fig. 2

*Holotype.* Plate 48, figs. 2a, b. Specimen Sy80/2/1/B; Sylvan Shale, 80 ft. (24.38 m.) stratigraphically above base, Sycamore Creek.

*Diagnosis.* Small conical to pyriform test. Maximum diameter about four-fifths total length; base convex, margin rounded. Neck weakly differentiated from chamber, short, tapering; commonly not developed. Aperture equal to, or slightly greater than, half maximum diameter. Numerous closely spaced slender processes; arms of adjacent

processes united, forming a complete reticulum that stands as high as  $12\ \mu$  above surface of test wall.

*Dimensions (in microns)*. 25 specimens measured.

	Total length	Maximum diameter	Apertural diameter
Holotype:	128	89	53
Range:	105–158	81–106	40–63
Mean:	123	93	51

*Remarks.* *Acanthochitina rashidi* is considered as a separate species because it falls well outside Jansonius's (1964, p. 909) definition of *Kalochitina multispinata*. Wherever it has been found, however, *A. rashidi* makes up continuously intergrading populations with *K. multispinata* and many transitional forms occur. The two species are distinguished arbitrarily and solely on the basis of their ornaments. Their close relationship is not reflected in the strictly empirical system of classification followed here, however, and they are unavoidably referred to different genera.



TEXT-FIG. 2. *Acanthochitina rashidi* sp. nov. Lateral profile,  $\times 400$ . For the sake of clarity only the ornament seen in profile is shown.

*Comparison.* In *Hercochitina downiei* Jenkins 1967 processes stand in longitudinal rows, and most frequently are connected only to those longitudinally adjacent to them. *A. rashidi* is much smaller than the type species (total length of 25 British examples  $300\text{--}485\ \mu$ , mean  $408\ \mu$ ), and has a much more delicate ornament; processes on the basal margin do not differ appreciably from those elsewhere and, in particular, are not connected by a membrane.


*Material.* Several thousand single tests, and a dozen chains of two or three tests each.

*Occurrence.* Sy80–Sy200. *A. rashidi* is restricted to the middle of the Sylvan Shale, although forms transitional to it, but referable to *K. multispinata*, occur at the younger horizons Sy220, Sy240, and Sy260.

#### EXPLANATION OF PLATE 47

Figs. 1–19. *Ancyrochitina merga* sp. nov., all but 19  $\times 250$ . 1–8, Sy80/1/1/B, Sy60/1/1/F, B, E, Sy80/1/1/A, D, C, and Sy60/1/1/H, respectively, bearing a variety of T- and Y-shaped appendices. 9–11, Sy60/1/1/D, Sy80/1/1/E and Sy60/1/1/A, respectively, with  $\lambda$ -shaped appendices. 12, 13, Sy80/1/1/F (holotype) and Sy60/1/1/C, respectively, each possessing T-, Y-, and  $\lambda$ -shaped appendices. 14–17, Sy40/1/1/A, C, B, and D, respectively, with ornaments of simple spines and  $\lambda$ -spines; this form represents the species near the base of the Sylvan Shale, whereas younger Sylvan strata contain only smooth-walled forms (figs. 1–13). 18, Sy60/1/1/G, cluster of three tests; two are in polar view, the upper right in lateral view. 19, Sy80/1/1/G, detail showing three simple appendices, and one with three orders of branching, in phase-contrast illumination  $\times 400$ .

Fig. 20. *Acanthochitina rashidi* sp. nov., Sy100/2/1/B, in phase-contrast illumination,  $\times 400$ .



JENKINS, Ordovician chitinozoa from Oklahoma



Genus *ANCYROCHITINA* Eisenack 1955a

Type species. *Conochitina ancyrea* Eisenack 1931 (by original designation), Silurian, Baltic.

*Ancyrochitina merga* sp. nov.

Plate 47, figs. 1-19; text-fig. 3

*Holotype*. Plate 47, fig. 12. Specimen Sy80/1/1/F; Sylvan Shale, 80 ft. (24-38 m.) stratigraphically above base, Sycamore Creek.

*Diagnosis*. Small fungiform test. Chamber about half total length, wider than long; base slightly convex, margin rounded. 8-24, generally fewer than 15, simple or branching appendices up to one-third of the maximum diameter in length; generally 1-3, rarely 4, orders of Y- or T-shaped branching into 2 sharply diverging, equal distal limbs. Commonly, appendices divide ( $\lambda$ -shaped branching) in a transverse plane to form two proximal limbs which meet the basal margin abruptly. Oral tube cylindrical or slightly flaring, one-third to half maximum diameter in width; aperture fringed with hairs up to  $5\mu$  in length. Test smooth, or bearing slender, tapering, simple or  $\lambda$ -spines with pointed tips, up to one-sixth maximum diameter in length; thinly distributed, absent on base and basal margin.



TEXT-FIG. 3. *Ancyrochitina merga* sp. nov. Lateral view of test showing the appendices and the ornament of  $\lambda$ -spines and simple spines,  $\times 400$ .

*Dimensions (in microns)*. 25 specimens measured.

	Total length	Chamber length	Maximum diameter	Oral tube diameter	Apertural diameter	Appendix length
Holotype:	125	70	91	35	c. 35	< 30
Range:	100-155	52-71	70-98	32-46	35-55	< 35
Mean:	128	64	81	36	44	—

*Description*. The flanks of the short, wide chamber are almost straight and taper rapidly, but with no abrupt change of curvature, into the neck. The maximum diameter generally is 120-140% of the chamber length. Where *A. merga* first occurs (horizon Sy40) it bears slender, simple spines and  $\lambda$ -spines up to  $12\mu$  in length, distributed thinly over the test (Pl. 47, figs. 14-17). Throughout the remainder of the formation, however, forms with smooth walls, or forms with very few short spines ( $< 2\mu$  in length) on the shoulder and neck, greatly predominate over the more ornate forms.

*Comparison*. *A. merga* is readily distinguished by the style and number of its appendices. *A. corniculans* Jenkins 1969 from the Viola and Fernvale Limestones is cylindroconical rather than fungiform, and has only 4-6 appendices.

*Material*. Several thousand single tests.

*Occurrence*. Sy40-Sy305.

## Genus CLATHROCHITINA Eisenack 1959

*Type species. Clathrochitina clathrata* Eisenack 1959 (by original designation), Wenlock, Gotland.

*Diagnosis.* 'Chitinozoans shaped like *Ancyrochitina ancyrea*, and furnished with appendices whose distal ends coalesce with a ring situated concentrically about the basal margin' (Eisenack 1959, p. 15).

*Remarks.* *Clathrochitina* Eisenack 1959 was conceived for cylindroconical chitinozoans possessing *distinct appendices* (generally discrete processes confined to the basal margin) that coalesce distally; *Plectochitina* Cramer 1964 is a junior synonym. Unfortunately, several species lacking distinct appendices have been referred inappropriately to *Clathrochitina* on the basis of *finely meshed networks* suspended from their basal margins, and *Plectochitina* has assumed the taxonomic role originally intended for *Clathrochitina*. Consequently, the genus *Sagenachitina* (q.v., p. 270) is here proposed for chitinozoans with *networks* suspended from their basal margins; and several species are transferred to it from *Clathrochitina*. The latter is considered to include, besides the type species, *Clathrochitina multiramosa* (Taugourdeau and Jekhowsky 1960) Taugourdeau 1967, *C. saharica* (Taugourdeau 1962) comb. nov., *C. carminae* (Cramer 1964) comb. nov., *C. rosendae* (Cramer 1964) comb. nov., and *C. combazi* (Cramer 1967a) comb. nov. It seems essentially a Silurian genus, being known only from the uppermost Ordovician (this paper), the Silurian (Eisenack 1959; Taugourdeau 1962, 1967; Cramer 1964, 1967a; Taugourdeau *et al.* 1967, p. 85, under *Plectochitina*), and, perhaps, the basal Devonian (Cramer 1964).

Cramer (1967a, pp. 84, 94; 1967b, p. 47) interpreted the structure on the basal margin of *C. clathrata* (type species of its genus) as a 'perforate cingulum' and, on the basis of this interpretation, transferred the species to *Cyathochitina* Eisenack 1955b. For species previously referred to *Clathrochitina*, which was abandoned through the loss of its type species, he proposed the genera *Clathrochitinella* and *Pseudoclathrochitina*. While Cramer's proposals may seem reasonable in the light of his structural interpretation, the latter is entirely inconsistent with Eisenack's (1959, p. 15, pl. 1, figs. 3, 4; text-fig. 4) clear illustrations and lucid, unambiguous description of the genus and its type species. Cramer's proposals and transfer of *C. clathrata* are, if only for this reason, unacceptable, and *Clathrochitinella* and *Pseudoclathrochitina* should be abandoned along with *Plectochitina*. Aside from the faulty interpretive basis upon which they rest, Cramer's proposals and transfer serve little more than to complicate and confuse a relatively straightforward issue, namely the generic placement of two groups of cylindroconical chitinozoans, one characterized by networks suspended from the basal margin (*Sagenachitina*), the other by appendices which coalesce distally (*Clathrochitina*). The existence of a few forms transitional between *Sagenachitina* and *Clathrochitina* is known (e.g. Taugourdeau 1967, pl. 1, fig. 9) but scarcely complicates the issue.


*Clathrochitina sylvanica* sp. nov.

Plate 48, figs. 3-13; text-fig. 4

*Holotype.* Plate 48, figs. 11a, b. Specimen Da50/12/1/A; Sylvan Shale, 50 ft. (15-24 m.) stratigraphically above base, in outcrop on west side of U.S. Highway 77, about 4 miles (6.4 km.) south of Davis, Oklahoma.



*Diagnosis.* Conical chamber slightly longer than oral tube, approximately as wide as long; base flat or slightly convex, margin rounded. Appendices of uniform thickness and texture, suspended at 6–15, generally 8–12, points on the basal margin; commonly anastomosing; occasionally discrete for their full lengths and connected at their tips by a continuous ring; equal to, or less than, maximum diameter in length. Oral tube cylindrical, half maximum diameter in width. Test wall smooth.



TEXT-FIG. 4. *Clathrochitina sylvanica* sp. nov. Lateral views of two variants with strongly developed appendices,  $\times 400$ .

*Dimensions (in microns)*, 25 specimens measured.

	Total length	Chamber length	Maximum diameter	Oral tube diameter	Appendix length
Holotype:	120	62	78	36	< 53
Range:	110–160	60–85	75–103	36–48	12–90
Mean:	131	76	84	40	—

*Description.* The flanks taper fairly uniformly, and the junction of the chamber and neck generally is clearly defined. The appendices are very strongly developed. Their pattern of branching and anastomosing varies considerably at each horizon, but the same, or closely similar, variants occur wherever the species has been found.

*Comparison.* *Ancyrochitina merga* is readily distinguished from *C. sylvanica* by its discrete, pitchfork-shaped appendices. Damaged specimens which have lost their appendices, however, may be exceedingly difficult to identify, but typical examples of *A. merga* are, nevertheless, fungiform rather than cylindroconical, and may bear

simple or  $\lambda$ -spines. Four similar species of *Clathrochitina* differ from *C. sylvanica* as follows. *C. combazi* (Cramer 1967a) possesses 12–24 modestly branching appendices, whereas *C. roseadae* (Cramer 1964) bears only a few. The appendices of *C. clathrata* Eisenack 1959 are very short; those of *C. carminae* (Cramer 1964) branch elaborately and may extend as far as 130  $\mu$  from the basal margin. *C. multiramosa* (Taugourdeau and Jekhowsky 1960) Taugourdeau 1967 is readily distinguished by about 12 short processes attached immediately below the aperture, and connected at their tips by a continuous ring.

*Material.* Approximately 400 single tests.

*Occurrence.* ?Syl, Sy16–Sy180. *Clathrochitina sylvanica* is a distinctive element in approximately the lower half of the Sylvan Shale.

#### Genus SAGENACHITINA gen. nov.

*Type species.* *Clathrochitina oblonga* Benoit and Taugourdeau 1961 (by original designation), Arenig, Algeria.

*Diagnosis.* Chitinozoa with cylindroconical or campanulate tests and a network suspended from the basal margin.

*Remarks.* *Sagenachitina* gen. nov. is not represented in the Sylvan Shale, but is proposed for a group of closely similar species including 5 formerly referred inappropriately to *Clathrochitina* Eisenack 1959. These are *Sagenachitina oblonga* (Benoit and Taugourdeau 1961) comb. nov., *S. aquitanica* (Taugourdeau 1961) comb. nov., *S. eisenacki* (Taugourdeau 1961) comb. nov., *S. retifera* (Taugourdeau and Jekhowsky 1960) comb. nov., and *S. striata* (Benoit and Taugourdeau 1961) comb. nov. *Sagenachitina* differs from *Clathrochitina* (q.v., p. 268) in that its basal margin supports a finely divided network, rather than a relatively small number of distinct processes which coalesce distally. Apparently it is restricted to the Ordovician (Taugourdeau *et al.* 1967, p. 78), whereas *Clathrochitina* (*sensu* Eisenack 1959) seems essentially a Silurian genus.

*Acknowledgement.* *Sagenachitina* was introduced to the literature by Jansonius (1967, p. 352) as an informal manuscript name. It is validated here, in its original sense, with the permission and approval of Dr. Jansonius.

#### Genus CONOCHITINA Eisenack 1931 restr. 1955b

*Type species.* *Conochitina claviformis* Eisenack 1931 (by original designation), Silurian, Baltic.

#### *Conochitina elegans* Eisenack 1931

Plate 49, figs. 1–17

#### EXPLANATION OF PLATE 48

- Figs. 1, 2. *Acanthochitina rashidi* sp. nov.,  $\times 400$ . 1, Sy100/2/1/D, in phase-contrast illumination. 2, Sy80/2/1/B, holotype; in bright-field (2a) and phase-contrast (2b) illumination.
- Figs. 3–13. *Clathrochitina sylvanica* sp. nov., illustrating variation in the size and complexity of the appendices,  $\times 250$ . 3, Sy60/12/1/D. 4, Sy100/12/1/A. 5, Sy80/12/1/B. 6, Sy60/12/1/C, polar view. 7, Sy60/12/1/E. 8, Sy60/12/1/K. 9, Sy80/12/1/A. 10, Sy100/12/1/C. 11, Da50/12/1/A, holotype; in bright-field (11a) and phase-contrast (11b) illumination. 12, Sy60/12/1/H. 13, Sy60/12/1/A.



1




2a



2b



3



4



5



6



7



8



9



10



11a



11b



12



13

JENKINS, Ordovician chitinozoa from Oklahoma



- 1931 *Conochitina elegans* Eisenack, p. 87, pl. 2, fig. 4 (holotype).  
 1934 *Rhabdochitina conocephala* Eisenack, p. 61, pl. 4, figs. 10-12; text-fig. 32.  
 1959 *Conochitina elegans* Eisenack; Eisenack, p. 3, pl. 2, figs. 4 (neotype), 5; text-fig. 1.  
 1960 *Rhabdochitina conocephala* Eisenack; Taugourdeau and Jekhowsky, p. 1230, pl. 9, fig. 131.  
 non 1962 *Conochitina elegans* Eisenack; Beju and Dăneț, p. 531, pl. 1, figs. 31, 32.  
 non 1964 *Rhabdochitina conocephala* Eisenack; Cramer, p. 351, pl. 22, fig. 14; pl. 23, figs. 7, 11, 12.  
 1965 *Rhabdochitina hedlundii* Taugourdeau, p. 472, pl. 3, figs. 60, 66.  
 1965 *Conochitina elegans* Eisenack; Eisenack, p. 126, pl. 10, fig. 9.  
 1967 *Conochitina elegans* Eisenack; Jenkins 1967, p. 455, pl. 71, figs. 1-4.

*Dimensions (in microns). 50 specimens measured.*

		Total length	Maximum diameter	Minimum diameter	Apertural diameter
	Range:	204-904	62-96	50-78	61-78
	Mean:	403	78	61	69
26 specimens from Estonia (Eisenack 1959)	Range:	288-667	—	—	—
	Mean:	467	—	—	—
26 specimens from the Ostseekalk of south Finland (Eisenack 1965)	Range:	332-690	—	—	—
	Mean:	493	—	—	—
30 specimens from England (Jenkins 1967)	Range:	200-616	58-92	—	—
	Mean:	388	73	—	—

*Remarks.* *Conochitina elegans* Eisenack 1931 shows the same pattern of morphological variation in the Sylvan Shale as it does in the Caradocian Jewe and Kegel Beds, D<sub>1</sub>-D<sub>2</sub>, of Estonia (Eisenack 1959, p. 3, text-fig. 1), and in the Caradoc Series of England (Jenkins 1967). Throughout the Sylvan Shale, short conical or cylindroconical forms up to 400  $\mu$  in length (Pl. 49, figs. 8-15) predominate numerically over much longer cylindrical forms, up to 900  $\mu$  in length, which have pronounced aboral swellings (Pl. 49, figs. 1-7). The aperture is distinctively fringed by numerous, irregular, simple or branching processes up to 6  $\mu$  in length.

I am reasonably sure that *Conochitina elegans* and *Rhabdochitina hedlundii* Taugourdeau 1965 are conspecific. Well-preserved typical examples of *C. elegans* occur at several horizons in the Upper Ordovician Maquoketa Formation from Iowa, where Taugourdeau obtained the type material of *R. hedlundii*, but no other species comparable to *C. elegans* or *R. hedlundii* is represented.


*Material.* Several thousand single tests, and clusters of up to 15 tests each.

*Occurrence.* Syl-Sy305. The species occurs in the Caradoc Series of Estonia (Eisenack 1959, 1962b) and Shropshire, England (Jenkins 1967), and in the Ostseekalk of north Germany and south Finland (Eisenack 1965). Taugourdeau and Jekhowsky (1960) record it, as *Rhabdochitina conocephala*, in Algerian sediments referred with reservations (op. cit., pp. 1205, 1230) to the lower Silurian. Eisenack (1964) refers to *C. cf. elegans* a group of forms from the Llandovery, Wenlock, and lower Ludlow of Gotland which, while closely similar to the Ordovician type material, are not certainly conspecific with it.

*Conochitina cactacea* Eisenack 1937

Plate 49, figs. 18–25; text-fig. 5

- 1937 *Conochitina cactacea* Eisenack, p. 222, pl. 15, figs. 14, 15 (holotype).  
 1959 *Conochitina cactacea* Eisenack; Eisenack, p. 10, pl. 1, figs. 12 (neotype), 13; text-fig. 2a, b.  
 non 1965 *Conochitina* cf. *cactacea* Eisenack *typica* Taugourdeau, p. 467, pl. 1, fig. 10.  
 1965 *Conochitina cactacea* Eisenack; Eisenack, p. 125, pl. 9, figs. 18, 19.  
 1967 *Conochitina cactacea* Eisenack; Laufeld, p. 299, fig. 9.



TEXT-FIG. 5. *Conochitina cactacea* Eisenack 1937. Lateral views illustrating (left) a typical cylindroconical example with randomly distributed spines, and (right) a pyriform test whose spines are expanded proximally and arranged in longitudinal rows,  $\times 350$ .

Dimensions (in microns), 25 specimens measured.

	Total length	Maximum diameter	Oral tube diameter	Apertural diameter	Spine length
Range:	118–240	80–123	52–77	53–73	< 15
Mean:	165	97	63	62	—
Neotype (Eisenack 1959):	123	72	42	43	—

## EXPLANATION OF PLATE 49

Figs. 1–17. *Conochitina elegans* Eisenack 1931,  $\times 100$ . 1–7, Sy60/4/1/A, D, K, Da3/4/1/B, Sy60/4/1/R, G, and Sy100/4/1/C, respectively, long cylindrical tests with conspicuous aboral swellings. 8–10, Sy60/4/1/S, H, and J, respectively, shorter tests with neck and chamber differentiated. 11–17, Sy40/4/1/C, D, F, A, Sy60/4/1/T, Da3/4/1/D and Sy60/4/1/P, respectively, short to very short tests illustrating some of the very considerable variation that is expressed in both the large and small size-fractions of populations of *C. elegans*.

Figs. 18–25. *Conochitina cactacea* Eisenack 1937, all except 19b  $\times 250$ . The figured specimens have been chosen to illustrate the ornament; most are distorted and none shows the typical shape of this well-known species. 18, Sy60/6/1/A. 19, Sy16/6/1/A; 19a, showing exceptionally well-preserved ornament in upper right quadrant of photograph; 19b, detail of proximally-expanded spines, in phase-contrast illumination  $\times 1600$ . 20, Sy16/6/1/D. 21, Sy16/6/1/B. 22, Sy16/6/1/F. 23, Sy60/6/1/B. 24, Sy16/6/1/E. 25, Sy16/6/1/C.





JENKINS, Ordovician chitinozoa from Oklahoma



*Description.* The chamber is conical with straight or swollen flanks, and makes up more than half the total length. The base is flat, with a broadly rounded margin. The oral tube, which occasionally is not developed, is cylindrical and approximately two-thirds of the maximum diameter in width. The aperture is straight, or fringed with irregularly spaced processes up to  $3\ \mu$  in length. Short spines, most of which are simple, cover the entire test. Branching spines with two distal limbs, and  $\lambda$ -spines with two or three discrete bases, however, are common. Spines may taper uniformly, or their proximal portions may be widely expanded in a plane parallel with the test's longitudinal axis. Commonly the spines show some tendency to loosely align themselves in longitudinal rows, and in some examples the expanded proximal portions of longitudinally adjacent spines occasionally coalesce, forming very short ridges up to  $3\ \mu$  in height. Such ridges tend to be less strongly developed, though no less common, toward the aperture. In general, spines are largest on the basal margin and become smaller orally.

*Material.* Approximately 200 single tests.

*Occurrence.* Syl-Sy305. *C. cactacea* occurs throughout the Sylvan Shale, but nowhere is it common. The species occurs in the Ostseekalk of north Germany and Gotland (Eisenack 1965), and in the Caradocian Dalby and Slandrom Formations of central Sweden (Laufeld 1967). A few atypical examples are known from the Wesenberg Beds, E, of Estonia (Eisenack 1962b, 1965).

#### Genus CYATHOCHITINA Eisenack 1955b

*Type species.* *Conochitina campanulaeformis* Eisenack 1931 (by original designation), Ordovician, Baltic.

#### *Cyathochitina agrestis* sp. nov.

Plate 50, figs. 11, 18

*Holotype.* Plate 50, figs. 11a, b. Specimen Syl/8/1/A; Sylvan Shale, 1 ft. (0.3 m.) stratigraphically above base, Sycamore Creek.

*Diagnosis.* Large test. Chamber conical to swollen cylindrical, about two-thirds total length; maximum diameter in lower half or middle of chamber, about one-third chamber length; base flat, about three-quarters maximum diameter in width. Carina attached some distance aborally of maximum diameter. Oral tube cylindrical or slightly flaring, about four-fifths maximum diameter in width; aperture straight. Test wall rough.

*Dimensions (in microns).* 6 specimens measured.

	Total length	Chamber length	Maximum diameter	Oral tube diameter	Apertural diameter	Carina width
Holotype:	980	660	192	148	172	< 20
Range:	592-980	376-660	176-192	128-152	136-172	< 25
Mean:	784	549	184	142	153	—

*Comparison.* In general shape this species closely resembles *Acanthochitina barbata* Eisenack 1931 and *Cyathochitina stentor* (Eisenack 1937), but lacks the diagnostic ornament of either (Jenkins 1967, pp. 443-5; Laufeld 1967, p. 318, respectively). It is much larger than *Cyathochitina calix* (Eisenack 1931), 57 Baltic examples of which (Eisenack 1962a) average  $299\ \mu$  in total length (min.  $190\ \mu$ , max.  $450\ \mu$ ).

*Material.* 10 single tests.

*Occurrence.* Syl.

*Cyathochitina ontariensis* (Jansonius 1964) comb. nov. emend.

Plate 50, figs. 1-9

1964 *Tanuchitina ontariensis* Jansonius, p. 910, pl. 1, figs. 5, 6 (holotype).

*Emended diagnosis.* Elongate, cylindroconical, or campanulate test. Chamber approximately two-thirds total length; base almost flat; margin drawn out into a short sharp carina, generally directed aborally, and situated slightly below the maximum diameter. Neck cylindrical, half to two-thirds maximum diameter in width; collarette flaring; aperture serrate or fimbriate, generally 8-12  $\mu$  wider than neck. Wall smooth.

*Dimensions (in microns).* 25 specimens measured.

	Total length	Maximum diameter	Neck diameter	Apertural diameter	Carina width
Range:	268-740	102-180	56-88	64-106	< 3
Mean:	475	142	69	79	—
Holotype (Jansonius 1964):	310	—	—	—	—

*Description.* The flanks may taper uniformly or be slightly swollen (Pl. 50, figs. 1, 2, 4, 5), in which cases the chamber and neck are more or less clearly distinguishable. Occasionally the test is trumpet-shaped, having concave flanks which merge with the neck (Pl. 50, figs. 3, 7). Originally the base probably was flat or almost so; in most specimens, however, perhaps owing to compression, it is convex in the centre and concave toward the margin. The carina is a continuous, uniformly wide, knife-edge rim which, in lateral view, rarely protrudes more than 3  $\mu$  beyond the general silhouette of the test.

*Remarks.* This species was designated type species of *Tanuchitina* by Jansonius (1964), but is here considered to fall well within the scope of *Cyathochitina* Eisenack 1955b.


*Comparison.* *C. agrestis* sp. nov. has a much larger carina than *C. ontariensis* and the shape of its test is quite different. *C. stentor* (Eisenack 1937) is furnished with a strongly developed, skirt-like carina (up to 85  $\mu$  in width; Laufeld 1967, p. 318) and an ornament of strongly defined longitudinal ridges.

*Material.* Several thousand single tests.

*Occurrence.* Sy80-Sy305. The type material is from the subsurface Upper Ordovician of Ontario.

## EXPLANATION OF PLATE 50

- Figs. 1-9. *Cyathochitina ontariensis* Jansonius 1964,  $\times 100$ . 1, 2, Sy100/7/1/A and B, respectively, typical examples with slender campanulate tests and weakly developed carinae. 3, Sy80/7/1/B, trumpet-shaped test. 4, 5, Sy80/7/1/A and Sy100/70/1/C, respectively, with well-developed carinae. 6-7, Sy80/7/1/C and Sy100/7/1/D, respectively, very short tests. 8, Sy80/7/1/E. 9, Sy80/7/1/D. Figs. 10, 12-17, 19, 20. *Desmochitina minor* Eisenack 1931. 10, 14, 15, 19, 20, Sy100/9/1/A, Sy80/9/1/A, B, C, and Sy100/9/1/B, single tests each with an operculum firmly attached at the base,  $\times 250$ . 12, Sy40/9/1/A, single test lacking an operculum at its base (presumably one was once present and has since been lost), but having retained the operculum sealing its own aperture (this is set squarely in the throat of the specimen),  $\times 250$ . 13, 16, 17, Sy60/9/1/B, H, and Sy100/9/1/C, respectively, chains of tests connected aperture-to-base,  $\times 100$ . Figs. 11, 18. *Cyathochitina agrestis* sp. nov. 11, Syl/8/1/A, holotype; 11a,  $\times 100$ ; 11b, carina,  $\times 250$ . 18, Syl/8/1/B,  $\times 100$ .
- Fig. 21. *Desmochitina scabiosa* (Wilson and Hedlund 1964), Sy60/10/1/D, cluster of four tests,  $\times 250$ .



JENKINS, Ordovician chitinozoa from Oklahoma





Genus *DESMOCHITINA* Eisenack 1931 emend. 1962a

*Type species. Desmochitina nodosa* Eisenack 1931 (by original designation), Ordovician, Baltic.

*Remarks.* For the present I share Eisenack's (1968, pp. 155, 185) view that *Hoegisphaera* Staplin 1961 is superfluous, and would place it (and its junior synonym *Calpichitina* Wilson and Hedlund 1964) in synonymy with *Desmochitina* Eisenack 1931 emend. 1962a. *Hoegisphaera* was established by Staplin (1961) for 'a new type of Paleozoic microfossil possibly allied to the Chitinozoa' (op. cit., p. 392), in which 'The colour and texture of the wall are similar to those of Chitinozoa, but the analogy cannot be carried farther' (op. cit., p. 419). However, it is clear that the species assigned to *Hoegisphaera* are chitinozoans, and they have been widely recognized as such in the literature (Jansonius 1964, p. 913; 1967, p. 350; Taugourdeau *et al.* 1967, p. 61; Laufeld 1967, pp. 319-20, 327-8; Jenkins 1967, p. 462; Eisenack 1968, pp. 155, 185). Three years after *Hoegisphaera* had been established, Wilson and Hedlund (1964) described the genus *Calpichitina* with its type species *C. scabiosa*. This generic name, however, was proposed for a species (*C. scabiosa*) which clearly fell within Staplin's (1961) concept of *Hoegisphaera*, and was closely similar in size and shape to *Desmochitina conplanata* Eisenack 1932 (for dimensions see Eisenack 1959, p. 16). Very shortly after the establishment of *Calpichitina*, Wilson and Dolly (1964), doubting its validity, abandoned it by transferring the type species (*C. scabiosa*) to *Hoegisphaera*.

*Desmochitina minor* Eisenack 1931

Plate 50, figs. 10, 12-17, 19, 20

1969 *Desmochitina minor* Eisenack; Jenkins, pp. 20, 21, pl. 6, figs. 1-18 (q.v. for further synonymy).

*Dimensions (in microns).* 25 specimens measured.

	Total length	Chamber length	Maximum diameter	Minimum (neck) diameter	Apertural diameter
Range:	72-120	60-108	66-86	46-60	54-64
Mean:	92	84	76	53	58

*Remarks.* Eight informal infraspecific taxa have been referred to *Desmochitina minor* by Eisenack (1958, 1962a). The form recorded here corresponds exactly to the informal taxon *D. minor* forma *typica* Eisenack 1958 but is considered to merit recognition as a separate species. It occurs widely within North America and Europe, where it is generally quite distinct. The remaining seven infraspecific taxa, and forms transitional to them, are apparently lacking in the Sylvan Shale.

Most specimens are indistinguishable from many of the examples from Bohemia (Eisenack 1948, text-figs. 14, 15), the Ordovician Rhenish Schiefergebirge (Eisenack 1939, text-figs. 1-3, 6), and the Swedish and British Caradoc (Laufeld 1967, fig. 25; Jenkins 1967, pl. 71, fig. 18). They differ appreciably, however, from Eisenack's Baltic material (1962a, pl. 16, figs. 1-8, 10; 1965, pl. 10, figs. 16, 17) and the Viola-Fernvale specimens recorded in an earlier study (Jenkins 1969), in that the general shape of the chamber in lateral view tends to be spherical rather than quadrangular; the oral tube is smaller and less sharply flaring; and chains of up to 20 tests are common.

In marked contrast to its broad pattern of morphological variation in the Viola and Fernvale Limestones (Jenkins 1969, p. 21, pl. 6, figs. 1-18), the species is represented

within the Sylvan Shale by a conservative form which, throughout the formation, varies little in size, shape, or surface texture.

*Material.* Many thousand single tests, and several hundred chains of up to 20 tests each. Clusters of up to 60 tests are common at horizon Sy60.

*Occurrence.* Sy16-Sy260. In the lower half of the formation (Sy16-Sy140) *D. minor* is a numerically important element in the chitinozoan fauna and occurs in long chains; thereafter its numbers are drastically reduced, and the species is missing or extremely scarce near the top of the formation (Sy280-Sy305). Its occurrence outside North America is given elsewhere (Jenkins 1969, p. 21).

*Desmochitina scabiosa* (Wilson and Hedlund 1964) Jenkins 1969

Plate 50, fig. 21

- 1958 *Desmochitina* sp. 2 Wilson, pl. 1, fig. 7.  
 1962a *Desmochitina* sp. Eisenack, p. 304, pl. 16, figs. 11, 12.  
 1964 *Calpichitina scabiosa* Wilson and Hedlund, p. 164, pl. 1, figs. 1 (holotype), 2-12.  
 1965 *Desmochitina lecaniella* Eisenack, p. 131, pl. 10, figs. 21, 22 (holotype).  
 1967 *Desmochitina lecaniella* Eisenack; Laufeld, p. 326, fig. 24.  
 1969 *Desmochitina scabiosa* (Wilson and Hedlund); Jenkins, p. 23.

*Dimensions (in microns).* 25 specimens measured.

	Total length	Chamber length	Maximum diameter	Apertural diameter
Range:	50-75	43-68	52-92	48-62
Mean:	61	56	73	56
Holotype (Wilson and Hedlund 1964):	60	—	79	41

*Remarks.* The stratigraphical ranges of *D. scabiosa* (Wilson and Hedlund 1964) and *D. lecaniella* Eisenack 1965 both lie within the Upper Ordovician, and the two species are so closely similar that it would seem impractical to continue distinguishing them. Populations from the Sylvan Shale consist largely of single tests, but clusters (*sensu* Kozlowski 1963) are common at several horizons.


*Material.* Several thousand single tests, and numerous clusters of up to 11 tests each.

*Occurrence.* Sy40-Sy305. Wilson and Hedlund (1964) illustrate several single tests particularly well, and record the species approximately 50 ft. (15.24 m.) above the base of the formation, at an outcrop on U.S. Highway 77 about 4 miles (6.4 km.) south of Davis, Oklahoma (text-fig. 1) and 20 miles (32 km.) north-west of the Sycamore Creek section. Taugourdeau records *C. scabiosa* (illustrated as *C. cf. scabiosa*) in the Maquoketa Formation, Richmond, of Iowa. *Desmochitina lecaniella* has been recorded in the Caradocian Kegel Beds, D<sub>2</sub>, of Estonia (Eisenack 1962a, 1965); in the Ostseekalk of north Germany, Gotland, and south Finland (Eisenack 1965); and in the Dalby Formation, Caradoc, of central Sweden (Laufeld 1967).

Genus KALOCHITINA Jansonius 1964

*Type species.* *Kalochitina multispinata* Jansonius 1964 (by original designation), Upper Ordovician, Ontario.

*Remarks.* The chief diagnostic features of this genus are its pyriform test, reduced neck, and ornament of numerous, generally small spines which frequently are aligned in longitudinal rows.



TEXT-FIG. 6. *Kalochitina multispinata* Jansonius 1964. Lateral views illustrating variation in the style of the ornament,  $\times 400$ . a, Typical example with evenly spaced spines. b, Typical example with spines showing tendency to align themselves in longitudinal rows. c, Example with reduced ornament of short thorn-like spines. d, Example furnished with ridges which appear to have formed, figuratively, as if by fusion of closely spaced  $\lambda$ -spines.

*Kalochitina multispinata* Jansonius 1964

Plate 51, figs. 1-10, 15; text-fig. 6

1958 Genus B, Wilson, pl. 1, figs. 10, 11.

1964 *Kalochitina multispinata* Jansonius, p. 909, pl. 2, figs. 21 (holotype), 22.

Dimensions (in microns). 30 specimens measured.

	Total length	Maximum diameter	Apertural diameter	Spine length
Range:	102-156	82-103	42-60	< 12
Mean:	121	90	51	—
Holotype (Jansonius 1964):	140	—	—	—

*Description.* The test is pyriform to conical, with a maximum diameter about four-fifths of the total length. The neck is short, tapering, and rather weakly differentiated from the chamber; in most individuals it has been lost or was never developed. The apertural diameter equals or slightly exceeds half the maximum diameter.

In all populations of *K. multispinata* the ornament varies considerably, but, for the most part, its range of variation is much the same throughout the formation. The lower and upper beds (up to horizon Sy60 and above horizon Sy200) contain pure populations of *K. multispinata*, but throughout approximately the middle 140 ft. (43 m.) of the formation (horizons Sy80 to Sy200) the species makes up continuously intergrading populations with *A. rashidi*. The two species are distinguished from each other arbitrarily and solely on the basis of their ornaments (compare text-figs. 2 and 6). Typical examples of *K. multispinata* are covered with closely and evenly spaced  $\lambda$ -spines, up to 12  $\mu$  in length and possessing 2-6 proximal limbs (text-fig. 6a). In many, but by no means all, of the more typical examples, the proximal limbs of each  $\lambda$ -spine, and the spines as a whole, show some tendency to loosely align themselves in approximately 30 more or less clearly defined longitudinal rows (text-fig. 6b). In addition, each population contains individuals whose ornament is reduced to short ( $< 2.5 \mu$  in length) thorn-like spines (text-fig. 6c). A rare form, seemingly of sporadic vertical distribution, is furnished with perforate or imperforate ridges up to 8  $\mu$  in height (text-fig. 6d), which appear to have formed, figuratively, as if by fusion of very closely spaced  $\lambda$ -spines. Variants transitional to *A. rashidi* (text-fig. 2) occur from horizon Sy80 to Sy260.

*Material.* Several thousand single tests, and 20 chains of up to 4 tests each.

*Occurrence.* Sy16-Sy260. The type material is from the subsurface Upper Ordovician of Ontario (Jansonius 1964).

### Genus RHABDOCHITINA Eisenack 1931

*Type species.* *Rhabdochitina magna* Eisenack 1931 (by original designation), Ordovician, Baltic.

#### *Rhabdochitina* sp.

Plate 51, fig. 14


*Remarks.* A species of very large chitinozoan is present in the Sylvan Shale, but only 2 incomplete examples are recorded here, one from horizon Sy80, the other from Sy120.

#### EXPLANATION OF PLATE 51

Figs. 1-10, 15. *Kalochitina multispinata* Jansonius 1964, all but 10  $\times 250$ . 1-5, Sy40/3/1/C, Sy100/3/1/A, E, G and Sy60/3/1/D, respectively. 6, Sy40/3/1/A, showing ornament through translucent test. 7, 8, Sy100/3/1/B and Sy40/3/1/D, respectively. 9, Da2/3/1/D, cluster of four tests showing a roughly parallel alignment of their longitudinal axes, a particularly common phenomenon in this species. 10, Sy40/3/1/M,  $\lambda$ -spines, in phase-contrast illumination  $\times 1000$ . 15, Sy100/3/1/D, chain of two tests.

Figs. 11-13, 16-20. *Sphaerochitina lepta* sp. nov., a series of tests illustrating *inter alia* the variable length of the neck,  $\times 250$ . 11, Sy60/11/1/F, compressed so the longitudinal axis of the neck has folded into the same plane as the transverse axis of the chamber (cf. text-fig. 7a). 12, 13, 16-18, Sy60/11/1/K (holotype), E, M, R, and B, respectively. These specimens are not carinate, as might be erroneously assumed from the photographs. The sharp prominences on the chambers are not original features but occur in tests where accommodation during compression has involved an inward collapse of the base (cf. text-fig. 7b). 19, 20, Sy100/11/1/C and D, with relatively long necks.

Fig. 14. *Rhabdochitina* sp., Sy80/5/1/A,  $\times 100$ .



JENKINS, Ordovician chitinozoa from Oklahoma

